



INTERNATIONAL ATOMIC ENERGY AGENCY

FOURTEENTH INTERNATIONAL CONFERENCE ON PLASMA
PHYSICS AND CONTROLLED NUCLEAR FUSION RESEARCH

Würzburg, Germany, 30 September – 7 October 1992

IAEA-CN-56/D-1-4

3D MHD Study of Helias and Heliotron


T. Hayashi, T. Sato, W. Lotz, P. Merkel, J. Nührenberg,
U. Schwenn and E. Strumberger

(Received – Aug. 21, 1992)

NIFS-168

Sep. 1992

AD-A257 116



DTIC
ELECTRONIC
NOV 1992
8
E

RESEARCH REPORT
NIFS Series

92-28953



This report was prepared for the International Atomic Energy Agency (IAEA) by the National Institute for Fusion Science (NIFS). It is intended for use in scientific meetings. Because of the provisional nature of its content, it is not to be used for official publication, the preprint is made available on the understanding that it may be reproduced in its present form. The views expressed and the conclusions drawn are those of the author(s) and do not necessarily reflect those of the government(s). In particular, neither the IAEA nor any other organization is responsible for any material reproduced in this preprint.

BEST
AVAILABLE COPY

NAGOYA, JAPAN

92 11 05 009

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.



INTERNATIONAL ATOMIC ENERGY AGENCY

**FOURTEENTH INTERNATIONAL CONFERENCE ON PLASMA
PHYSICS AND CONTROLLED NUCLEAR FUSION RESEARCH**

Würzburg, Germany, 30 September – 7 October 1992

IAEA-CN-56/D-1-4

3D MHD Study of Helias and Heliotron

T.HAYASHI, T.SATO

National Institute for Fusion Science

Furocho, Nagoya 464-01, Japan

DTIC QUALITY INSPECTED 4

W.LOTZ, P.MERKEL, J.NÜHRENBURG,

U.SCHWENN, E.STRUMBERGER

Max-Planck-Institut für Plasmaphysik

IPP-EURATOM Association

D-8046 Garching bei München

Federal Republic of Germany

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

key words: helical system, helias, heliotron, torsatron, stellarator, three dimensional equilibrium, beta limit, magnetic island, self-healing of magnetic island, divertor, finite pressure effect

This is a preprint or a paper intended for presentation at a scientific meeting. Because of the provisional nature of its content and since changes of substance or detail may have to be made before publication, the preprint is made available on the understanding that it will not be cited in the literature or in any way be reproduced in its present form. The views expressed and the statements made remain the responsibility of the named author(s); the views do not necessarily reflect those of the government of the designating Member State(s) or of the designating organization(s). In particular, neither the IAEA nor any other organization or body sponsoring this meeting can be held responsible for any material reproduced in this preprint.

3D MHD Study of Helias and Heliotron

ABSTRACT

Properties of magnetic islands induced by the finite pressure effect are numerically analyzed for three-dimensional magnetohydrostatic equilibria of Helias and $l = 2$ heliotron/torsatron types. For Helias, it is found that an island chain is generated on the $5/6$ rational surface, when such a surface appears in the plasma region of a finite- β equilibrium. The island chain, however, is not so dangerous as to destroy the plasma confinement even if it appears in a vanishingly small shear region. Moreover, it is definitely confirmed that the finite pressure effect sometimes exhibits a surprisingly good aspect, namely, that the vacuum islands are removed as β increases, which can be called 'self-healing' of islands. This property can be explained by the numerically discovered fact that the phases of islands induced by the finite pressure effect are always locked i.e. do not change regardless of β . The way islands appear at finite- β for heliotron/torsatron is significantly different from that for Helias. However, we find the self-healing of islands can also occur for the heliotron configuration. This phenomenon can partly explain the reason why the fragility of surfaces decreases when they are shifted inward by control of the external vertical field B_v , though the configuration develops a magnetic hill. Furthermore, we analyze the diversion properties of the magnetic field outside the last closed magnetic surface for finite- β equilibria, and find that divertor concepts which have been developed from the diversion properties of the corresponding vacuum fields can be maintained for finite- β equilibria for both configurations.

1. Introduction

Whether or not a three-dimensional (3D) magnetohydrostatic finite- β equilibrium can keep clearly nested magnetic surfaces is a long-pending question, but its physics has not yet fully been understood. Properties of magnetic islands induced by a finite pressure effect in typical helical systems, such as Helias proposed for W7-X and $l = 2$ heliotron/torsatron proposed for LHD, are quantitatively studied by a numerical method, namely the 3D equilibrium code HINT [1,2], which does not a priori demand the existence of regularly

nested magnetic surfaces. The equilibrium calculation is made for the half-pitch period of the configuration, keeping the stellarator symmetry.

In the process of equilibrium calculations, we find a remarkable property of magnetic islands induced by the finite pressure effect; the islands, in some cases, show the property of 'self-healing', i.e. the islands tend to shrink as β increases.

2. Formation of magnetic islands at finite- β in Helias

The magnetic surfaces of the W7-X configuration are designed in such a way that the rational surfaces, $\iota = 1, \frac{5}{6}$, can be avoided. It is however of interest to study the finite- β behaviour of the $n = 5/m = 6$ island chain to assess whether this would be dangerous or not. For this study a vacuum field situation was selected in which at finite- β this island chain occurs. The rotational transform at the magnetic axis, ι_0 , for the vacuum field is set slightly greater than $\frac{5}{6}$. One characteristic in the finite- β behaviour of the ι profile for Helias is that ι_0 decreases as β increases; in the case of $l = 2$ heliotron/torsatron configuration, ι_0 usually increases as a function of beta. Because of this behaviour, the $\frac{5}{6}$ rational surface can come into the plasma region; first it appears near the magnetic axis, and moves outward as beta increases. Thus, the $\frac{5}{6}$ island chain can appear when the $\frac{5}{6}$ surface resides at about half the plasma radius which is the position of vanishingly small shear. In this case the plasma β at the magnetic axis is $\beta_0 = 7.2\%$. As is shown in Fig.1, our results indicate that even this situation is not so dangerous as to destroy the plasma confinement. Most of the plasma region is covered by clearly nested surfaces. This is of course a consequence of the optimization of the configuration [3]. Corresponding to the formation of islands in the magnetic field structure as is shown in Fig.1 (a), the island structure is formed also in the pressure profile, as is shown in Fig.1 (b), where the pressure is almost flattened inside the $\frac{5}{6}$ island. We note that in the HINT code the pressure is automatically computed inside islands as a result of the incorporated calculation step which makes the pressure uniform along each field line. Coincidence of magnetic and pressure islands, which is for the first time demonstrated explicitly here in computational stellarator equilibria, proves validity of the code. Interestingly, local instabilities (Mercier and resistive interchange) are stabilized around the rational surface due to this flattening, the size of the islands being just large enough to bring about this stabilization.

When β is further increased, the $\frac{5}{6}$ island chain comes to the boundary region. Although

β is increased, the size of the island does not change so much, presumably due to the increased shear in the boundary region.

3. 'Self-healing' of magnetic islands in Helias

So far we have analyzed finite- β effects on equilibria from the point of view of formation of islands. Computational experience indicates that the phase induced by finite- β is independent of the value of β . Taking advantage of this property, we can have situations in which the finite- β effect acts rather to suppress islands which have existed in a vacuum field, if the phase of the vacuum field island is opposite to that of the β -induced ones. This process is confirmed in equilibrium calculations. A vacuum field is shown in Fig.2 (a), where a chain of magnetic islands is formed externally by controlling external currents, on the $\iota = \frac{5}{6}$ rational surface existing in the region of closed surfaces. Shown in Fig.2 (b) is the corresponding finite- β equilibrium with $\beta_0 = 9\%$. It is clearly observed that the $\frac{5}{6}$ island chain, which existed in the vacuum field, almost completely disappears, and a nice magnetic surface recovers by the effect of 'self-healing', though the $\iota = \frac{5}{6}$ surface still exists in the plasma region. Thus a high beta equilibrium keeping beautiful surfaces can be realized by making use of this feature.

When β is further increased, as is shown in Fig.2 (c) for $\beta_0 = 12\%$, the position of the $\iota = \frac{5}{6}$ surface moves further toward the boundary of the plasma, and the $\frac{5}{6}$ island chain appears again. Note, however, that the phase of the reappeared $\frac{5}{6}$ island chain is opposite to that of the vacuum one.

Based on the fact that islands may exhibit the property of self-healing described here, the physical equilibrium mechanism of formation of islands at rational surfaces in finite-pressure equilibria appears to be such that they are governed by resonant magnetic fields generated by the global effects of the plasma current density, mainly of the Pfirsch-Schlüter current, integrated over the whole plasma volume. We note that part of the reduction of the island size as β increases, shown in Fig.2 (b), may be attributed to the increase in the shear in the boundary region. However, the mere shear effect cannot explain the whole process shown here, since it does not explain the inversion in the phase of islands shown in Fig.2 (c). It appears natural to consider that the global plasma current effect, which determines the magnitude and the phase of the resonant field, mostly rules the island behaviour.

4. 'Self-healing' of magnetic islands in $l = 2$ heliotron/torsatron

We observe the nature of self-healing of islands not only in Helias but also in the $l = 2$ heliotron/torsatron configuration, which is characterized by medium to high shear. As is shown in Fig.3 (a) for typical vacuum surfaces of an $M = 10$ heliotron, the boundary region of the vacuum field is ergodic due to the loss of symmetry; in this case, two island chains, $n = 10/m = 8$ and $n = 10/m = 7$, are clearly visible near the outermost closed surface. It is a general property for vacuum fields of a wide range of $l = 2$ heliotron/torsatron configurations that the appearing islands are in phase at the outside of the torus when they are induced on several rational surfaces simultaneously [4]; here, both island chains have the x-point at the outside of the $\phi = \frac{\pi}{10}$ (horizontally elongated) poloidal cross section.

Because of the high shear nature, the way the islands appear at finite- β for heliotron/torsatron is significantly different from that for Helias; for heliotron, in general, the boundary region is ergodized due to overlapping of multiple island chains with smaller sizes and the boundary ergodized region gradually expands as β increases. However, we find the self-healing of islands can occur also for the heliotron configuration. Shown in Fig.3 (b) are magnetic surfaces for finite- β , $\beta_0 = 4.5\%$. When we observe the fine structure of boundary islands, it is visible that the $n = 10/m = 8$ island chain shrinks significantly, and the phase of the $n = 10/m = 7$ gets inverted. When the $n = 10/m = 7$ island chain disappears at slightly lower β , the minor radius of the outermost closed surface expands compared with that of the vacuum field. These island behaviours indicate that the finite pressure effect acts to generate resonant fields with a global structure with respect to the phase. In fact, when β further increases, island chains, all having the o-point at the outside of $\phi = \frac{\pi}{10}$, grow at several rational surfaces simultaneously, and eventually the boundary region is ergodized. In this way, the development of the expansion of the boundary ergodic region is delayed as β increases, because of the opposite nature in the phase of island chains between the vacuum field and the finite- β field for this case. This opposite nature in the phase typically appears when surfaces are shifted inward by control of the external vertical field B_v . This can partly explain the reason why fragility of surfaces at finite- β decreases as surfaces are shifted inward, although the configuration develops a magnetic hill.[1]

5. Divertor field structure in finite- β equilibria

One property of magnetic fields of helical systems is that divertor structures can be realized without installing any additional coils. We analyze the diversion properties of the magnetic field outside the last closed magnetic surface for finite- β equilibria, and investigate whether or not divertor concepts which have been developed from the diversion properties of the corresponding vacuum fields can be maintained for finite- β equilibria of heliotron and Helias, which has not been analyzed so far. For $l = 2$ heliotron/torsatron, the structure of magnetic surfaces near the plasma boundary for finite- β equilibria can be considerably different from that of the vacuum field with respect to the value of rotational transform or the extent of ergodic region. However, the position of the divertor field remains almost the same, which suggests that the same divertor plate can be used even in equilibria with quite high β . It should be noted that the 'width' of the divertor structure broadens when β increases. Therefore, one has to be careful in designing the divertor geometry, such as the baffle plate. For Helias, it is found that the finite- β equilibrium shows the same diversion properties as the vacuum field. Since the positions of the helical edges [5] are almost the same, the helical 'troughs' defined for the vacuum magnetic field work as well in the case of the finite- β equilibrium.

REFERENCES

- [1] HAYASHI, T., SATO, T., TAKEI, A., *Phys. Fluids B* 2 (1990) 329.
- [2] HAYASHI, T., TAKEI, A., SATO, T., *Phys. Fluids B* 4 (1992) 1539.
- [3] LOTZ, W., NÜHRENBURG, J., SCHWAB, C., IAEA-CN-53 (1990) Vienna 1991, II 603.
- [4] HAYASHI, T., TAKEI, A., OHYABU, N., SATO, T., *Nucl. Fusion* 31 (1991) 1767.
- [5] STRUMBERGER, E., *Nucl. Fusion* 32 (1992) 737.


Figure Captions

Fig.1 Magnetic surfaces (a) and pressure contours (b) of a finite- β Helias equilibrium selected for code validation: coincidence of magnetic and pressure islands, which is demonstrated here in computational stellarator equilibria for the first time.


Fig.2 'Self-healing' process of magnetic islands in a Helias equilibrium. (a) Vacuum magnetic surfaces which have a $5/6$ island chain. (b) Magnetic surfaces of a finite pressure equilibrium with $\beta_0 = 9\%$. The magnetic islands disappear and nice surfaces recover due to finite pressure effects, which is an opposite tendency compared to that found usually. (c) Magnetic surfaces of a higher β equilibrium with $\beta_0 = 12\%$. The island chain appears again, but with the inverted phase.

Fig.3 'Self-healing' process of magnetic islands in a $l = 2$ heliotron/torsatron equilibrium. (a) Vacuum magnetic surfaces which have $n = 10/m = 7$ and $n = 10/m = 8$ island chains in the boundary region. (b) Magnetic surfaces of a finite pressure equilibrium with $\beta_0 = 4.5\%$. The $n = 10/m = 8$ island chain almost disappears, and the phase of the $n = 10/m = 7$ gets inverted.

Helias



(a) Magnetic Field




(b) Pressure


Fig.1

'Self-healing' of magnetic islands

(a) Vacuum Field



(b) $\beta_0 = 9\%$



(c) $\beta_0 = 12\%$






Fig.2

$l = 2$ Heliotron/Torsatron



(a) Vacuum Field



(b) $\beta_0 = 4.5\%$

Fig.3

Recent Issues of NIFS Series

- NIFS-117 K. Itoh, H. Sanuki and S. -I. Itoh, *Fast Ion Loss and Radial Electric Field in Wendelstein VII-A Stellarator*; Oct. 1991
- NIFS-118 Y. Kondoh and Y. Hosaka, *Kernel Optimum Nearly-analytical Discretization (KOND) Method Applied to Parabolic Equations <<KOND-P Scheme>>*; Nov. 1991
- NIFS-119 T. Yabe and T. Ishikawa, *Two- and Three-Dimensional Simulation Code for Radiation-Hydrodynamics in ICF*; Nov. 1991
- NIFS-120 S. Kawata, M. Shiromoto and T. Teramoto, *Density-Carrying Particle Method for Fluid* ; Nov. 1991
- NIFS-121 T. Ishikawa, P. Y. Wang, K. Wakui and T. Yabe, *A Method for the High-speed Generation of Random Numbers with Arbitrary Distributions*; Nov. 1991
- NIFS-122 K. Yamazaki, H. Kaneko, Y. Taniguchi, O. Motojima and LHD Design Group, *Status of LHD Control System Design* ; Dec. 1991
- NIFS-123 Y. Kondoh, *Relaxed State of Energy in Incompressible Fluid and Incompressible MHD Fluid* ; Dec. 1991
- NIFS-124 K. Ida, S. Hidekuma, M. Kojima, Y. Miura, S. Tsuji, K. Hoshino, M. Mori, N. Suzuki, T. Yamauchi and JFT-2M Group, *Edge Poloidal Rotation Profiles of H-Mode Plasmas in the JFT-2M Tokamak* ; Dec. 1991
- NIFS-125 H. Sugama and M. Wakatani, *Statistical Analysis of Anomalous Transport in Resistive Interchange Turbulence* ;Dec. 1991
- NIFS-126 K. Narihara, *A Steady State Tokamak Operation by Use of Magnetic Monopoles* ; Dec. 1991
- NIFS-127 K. Itoh, S. -I. Itoh and A. Fukuyama, *Energy Transport in the Steady State Plasma Sustained by DC Helicity Current Drive* ;Jan. 1992
- NIFS-128 Y. Hamada, Y. Kawasumi, K. Masai, H. Iguchi, A. Fujisawa, JIPP T-IIU Group and Y. Abe, *New High Voltage Parallel Plate Analyzer* ; Jan. 1992
- NIFS-129 K. Ida and T. Kato, *Line-Emission Cross Sections for the Charge-exchange Reaction between Fully Stripped Carbon and Atomic Hydrogen in Tokamak Plasma*; Jan. 1992

- NIFS-130 T. Hayashi, A. Takei and T. Sato, *Magnetic Surface Breaking in 3D MHD Equilibria of $l=2$ Heliotron* ; Jan. 1992
- NIFS-131 K. Itoh, K. Ichiguchi and S. -I. Itoh, *Beta Limit of Resistive Plasma in Torsatron/Heliotron* ; Feb. 1992
- NIFS-132 K. Sato and F. Miyawaki, *Formation of Presheath and Current-Free Double Layer in a Two-Electron-Temperature Plasma* ; Feb. 1992
- NIFS-133 T. Maruyama and S. Kawata, *Superposed-Laser Electron Acceleration* Feb. 1992
- NIFS-134 Y. Miura, F. Okano, N. Suzuki, M. Mori, K. Hoshino, H. Maeda, T. Takizuka, JFT-2M Group, S.-I. Itoh and K. Itoh, *Rapid Change of Hydrogen Neutral Energy Distribution at L/H-Transition in JFT-2M H-mode* ; Feb. 1992
- NIFS-135 H. Ji, H. Toyama, A. Fujisawa, S. Shinohara and K. Miyamoto *Fluctuation and Edge Current Sustainment in a Reversed-Field-Pinch*; Feb. 1992
- NIFS-136 K. Sato and F. Miyawaki, *Heat Flow of a Two-Electron-Temperature Plasma through the Sheath in the Presence of Electron Emission*; Mar. 1992
- NIFS-137 T. Hayashi, U. Schwenn and E. Strumberger, *Field Line Diversion Properties of Finite β Helias Equilibria*; Mar. 1992
- NIFS-138 T. Yamagishi, *Kinetic Approach to Long Wave Length Modes in Rotating Plasmas*; Mar. 1992
- NIFS-139 K. Watanabe, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani, *Three-dimensional MHD Equilibrium in the Presence of Bootstrap Current for Large Helical Device (LHD)*; Mar. 1992
- NIFS-140 K. Itoh, S. -I. Itoh and A. Fukuyama, *Theory of Anomalous Transport in Toroidal Helical Plasmas*; Mar. 1992
- NIFS-141 Y. Kondoh, *Internal Structures of Self-Organized Relaxed States and Self-Similar Decay Phase*; Mar. 1992
- NIFS-142 U. Furukane, K. Sato, K. Takiyama and T. Oda, *Recombining Processes in a Cooling Plasma by Mixing of Initially Heated Gas*; Mar. 1992
- NIFS-143 Y. Hamada, K. Masai, Y. Kawasumi, H. Iguchi, A. Fijisawa and JIPP T-IIU Group, *New Method of Error Elimination in Potential Profile*

Measurement of Tokamak Plasmas by High Voltage Heavy Ion Beam Probes; Apr. 1992

- NIFS-144 N. Ohyabu, N. Noda, Hantao Ji, H. Akao, K. Akaishi, T. Ono, H. Kaneko, T. Kawamura, Y. Kubota, S. Morimoto, A. Sagara, T. Watanabe, K. Yamazaki and O. Motojima, *Helical Divertor in the Large Helical Device; May 1992*
- NIFS-145 K. Ohkubo and K. Matsumoto, *Coupling to the Lower Hybrid Waves with the Multijunction Grill; May 1992*
- NIFS-146 K. Itoh, S. -I. Itoh, A. Fukuyama, S. Tsuji and Allan J. Lichtenberg, *A Model of Major Disruption in Tokamaks; May 1992*
- NIFS-147 S. Sasaki, S. Takamura, M. Ueda, H. Iguchi, J. Fujita and K. Kadota, *Edge Plasma Density Reconstruction for Fast Monoenergetic Lithium Beam Probing; May 1992*
- NIFS-148 N. Nakajima, C. Z. Cheng and M. Okamoto, *High-n Helicity-induced Shear Alfvén Eigenmodes; May 1992*
- NIFS-149 A. Ando, Y. Takeiri, O. Kaneko, Y. Oka, M. Wada, and T. Kuroda, *Production of Negative Hydrogen Ions in a Large Multicusp Ion Source with Double-Magnetic Filter Configuration; May 1992*
- NIFS-150 N. Nakajima and M. Okamoto, *Effects of Fast Ions and an External Inductive Electric Field on the Neoclassical Parallel Flow, Current, and Rotation in General Toroidal Systems; May 1992*
- NIFS-151 Y. Takeiri, A. Ando, O. Kaneko, Y. Oka and T. Kuroda, *Negative Ion Extraction Characteristics of a Large Negative Ion Source with Double-Magnetic Filter Configuration; May 1992*
- NIFS-152 T. Tanabe, N. Noda and H. Nakamura, *Review of High Z Materials for PSI Applications ; Jun. 1992*
- NIFS-153 Sergey V. Bazdenkov and T. Sato, *On a Ballistic Method for Double Layer Regeneration in a Vlasov-Poisson Plasma; Jun. 1992*
- NIFS-154 J. Todoroki, *On the Lagrangian of the Linearized MHD Equations; Jun. 1992*
- NIFS-155 K. Sato, H. Katayama and F. Miyawaki, *Electrostatic Potential in a Collisionless Plasma Flow Along Open Magnetic Field Lines; Jun. 1992*
- NIFS-156 O.J.W.F.Kardaun, J.W.P.F.Kardaun, S.-I. Itoh and K. Itoh, *Discriminant Analysis of Plasma Fusion Data; Jun. 1992*

- NIFS-157 K. Itoh, S.-I. Itoh, A. Fukuyama and S. Tsuji, *Critical Issues and Experimental Examination on Sawtooth and Disruption Physics*; Jun. 1992
- NIFS-158 K. Itoh and S.-I. Itoh, *Transition to H-Mode by Energetic Electrons*; July 1992
- NIFS-159 K. Itoh, S.-I. Itoh and A. Fukuyama, *Steady State Tokamak Sustained by Bootstrap Current Without Seed Current*; July 1992
- NIFS-160 H. Sanuki, K. Itoh and S.-I. Itoh, *Effects of Nonclassical Ion Losses on Radial Electric Field in CHS Torsatron/Heliotron*; July 1992
- NIFS-161 O. Motojima, K. Akaishi, K. Fujii, S. Fujiwaka, S. Imagawa, H. Ji, H. Kaneko, S. Kitagawa, Y. Kubota, K. Matsuoka, T. Mito, S. Morimoto, A. Nishimura, K. Nishimura, N. Noda, I. Ohtake, N. Ohyabu, S. Okamura, A. Sagara, M. Sakamoto, S. Satoh, T. Satow, K. Takahata, H. Tamura, S. Tanahashi, T. Tsuzuki, S. Yamada, H. Yamada, K. Yamazaki, N. Yanagi, H. Yonezu, J. Yamamoto, M. Fujiwara and A. Iiyoshi, *Physics and Engineering Design Studies on Large Helical Device*; Aug. 1992
- NIFS-162 V. D. Pustovitov, *Refined Theory of Diamagnetic Effect in Stellarators*; Aug. 1992
- NIFS-163 K. Itoh, *A Review on Application of MHD Theory to Plasma Boundary Problems in Tokamaks*; Aug. 1992
- NIFS-164 Y. Kondoh and T. Sato, *Thought Analysis on Self-Organization Theories of MHD Plasma*; Aug. 1992
- NIFS-165 T. Seki, R. Kumazawa, T. Watari, M. Ono, Y. Yasaka, F. Shimpo, A. Ando, O. Kaneko, Y. Oka, K. Adati, R. Akiyama, Y. Hamada, S. Hidekuma, S. Hirokura, K. Ida, A. Karita, K. Kawahata, Y. Kawasumi, Y. Kitoh, T. Kohmoto, M. Kojima, K. Masai, S. Morita, K. Narihara, Y. Ogawa, K. Ohkubo, S. Okajima, T. Ozaki, M. Sakamoto, M. Sasao, K. Sato, K. N. Sato, H. Takahashi, Y. Taniguchi, K. Toi and T. Tsuzuki, *High Frequency Ion Bernstein Wave Heating Experiment on JIPP T-IIU Tokamak*; Aug. 1992
- NIFS-166 Vo Hong Anh and Nguyen Tien Dung, *A Synergetic Treatment of the Vortices Behaviour of a Plasma with Viscosity*; Sep. 1992
- NIFS-167 K. Watanabe and T. Sato, *A Triggering Mechanism of Fast Crash in Sawtooth Oscillation*; Sep. 1992
- NIFS-168 T. Hayashi, T. Sato, W. Lotz, P. Merkel, J. Nührenberg, U. Schwenn and E. Strumberger, *3D MHD Study of Helias and Heliotron*; Sep.